



Common fixed point results in extended b-metric spaces endowed with a directed graph

Cristian Chifu^a

^a*Babeş-Bolyai University Cluj-Napoca, Department of Business, Horea street, No.7, Cluj-Napoca, Romania*

Abstract

The purpose of this paper is to obtain some fixed point results in extended b-metric spaces for Reich-Rus and Ciric operators.

Keywords: extended b-metric spaces, fixed point, contraction, Reich-Rus and Ciric operators
2010 MSC: 47H10, 54H25.

1. Introduction

Definition 1.1. ([3]) Let X be a nonempty set and let $s \geq 1$ be a given real number. A functional $d : X \times X \rightarrow [0, \infty)$ is said to be a b -metric with constant s , if all axioms of the metric space take place with the following modification of the triangle axiom:

$$d(x, z) \leq s[d(x, y) + d(y, z)], \text{ for all } x, y, z \in X.$$

In this case the pair (X, d) is called a b -metric space with constant s .

Remark 1.2. The class of b -metric spaces is larger than the class of metric spaces since a b -metric space is a metric space when $s=1$. For more details and examples on b -metric spaces, see e.g. [2].

Example 1.3. Let $X = \mathbb{R}_+$ and $d : X \times X \rightarrow \mathbb{R}_+$ such that $d(x, y) = |x - y|^p$, $p > 1$. It's easy to see that d is a b -metric with $s = 2^p$, but is not a metric.

Email address: cristian.chifu@tbs.ubbcluj.ro (Cristian Chifu)

Definition 1.4. ([4]) Let X be a nonempty set and let $\theta : X \times X \rightarrow [1, \infty)$. A functional $d_\theta : X \times X \rightarrow [0, \infty)$ is said to be an extended b -metric, if for all $x, y, z \in X$, the following axioms are satisfied:

1. $d_\theta(x, y) = 0 \iff x = y$;
2. $d_\theta(x, y) = d_\theta(y, x)$;
3. $d_\theta(x, z) \leq \theta(x, y) [d_\theta(x, y) + d_\theta(y, z)]$, for all $x, y, z \in X$

In this case the pair (X, d_θ) is called an extended b -metric space.

Remark 1.5. If $\theta(x, y) = s$, for $s \geq 1$, then we obtain the definition of b -metric space.

Example 1.6. ([4]) Let $X = \{1, 2, 3\}$, $\theta : X \times X \rightarrow [1, \infty)$ and $d_\theta : X \times X \rightarrow [0, \infty)$ as:

$$\begin{aligned}\theta(x, y) &= 1 + x + y \\ d_\theta(1, 1) &= d_\theta(2, 2) = d_\theta(3, 3) = 0, \\ d_\theta(1, 2) &= d_\theta(2, 1) = 80, \\ d_\theta(1, 3) &= d_\theta(3, 1) = 1000, \\ d_\theta(2, 3) &= d_\theta(3, 2) = 600.\end{aligned}$$

Then d_θ is an extended b -metric.

Example 1.7. ([1]) Let $X = [0, 1]$, $\theta : X \times X \rightarrow [1, \infty)$ and $d_\theta : X \times X \rightarrow [0, \infty)$ as:

$$\begin{aligned}\theta(x, y) &= \frac{1 + x + y}{x + y} \\ d_\theta(x, y) &= \frac{1}{xy}, x, y \in (0, 1], x \neq y, \\ d_\theta(x, y) &= 0, x, y \in [0, 1], x = y, \\ d_\theta(x, 0) &= d_\theta(0, x) = \frac{1}{x}, x \in (0, 1].\end{aligned}$$

Then d_θ is an extended b -metric.

The following Lemma is very important in the proof of our results:

Lemma 1.8. ([1]) Let (X, d_θ) an extended b -metric space. If there exists $q \in (0, 1)$, such that the sequence $(x_n)_{n \in \mathbb{N}} \subset X$, for an arbitrary $x_0 \in X$, satisfies $\lim_{m, n \rightarrow \infty} \theta(x_m, x_n) < \frac{1}{q}$ and

$$0 < d_\theta(x_n, x_{n+1}) \leq qd_\theta(x_n, x_{n+1}), \text{ for any } n \in \mathbb{N}$$

then $(x_n)_{n \in \mathbb{N}}$ is a Cauchy sequence.

Let (X, d_θ) be an extended b -metric space and Δ be the diagonal of $X \times X$. Let G be a directed graph, such that the set $V(G)$ of its vertices coincides with X and $\Delta \subseteq E(G)$, where $E(G)$ is the set of the edges of the graph. Assume also that G has no parallel edges and, thus, one can identify G with the pair $(V(G), E(G))$.

Throughout the paper we shall say that G with the above mentioned properties *satisfies standard conditions*.

Let us denote by G^{-1} the graph obtained from G by reversing the direction of edges. Thus,

$$E(G^{-1}) = \{(x, y) \in X \times X : (y, x) \in E(G)\}.$$

Let us consider the mappings $T, S : X \rightarrow X$.

Definition 1.9. An element $x \in X$ is called *common fixed point* of the pair (T, S) , if $T(x) = S(x) = x$.

We shall denote by $CFix(T, S)$ the set of all common fixed points of the pair (T, S) , i.e.

$$CFix(T, S) = \{x \in X : T(x) = S(x) = x\}.$$

Definition 1.10. Suppose that $T, S : X \rightarrow X$ are two mappings on an extended b-metric space (X, d_θ) endowed with a directed graph G . We say that the pair (T, S) is *G-orbital-cyclic pair*, if for any $x \in X$,

$$\begin{aligned} (x, Tx) \in E(G) &\implies (Tx, STx) \in E(G) \\ (x, Sx) \in E(G) &\implies (Sx, TSx) \in E(G). \end{aligned}$$

Let us consider the following sets

$$\begin{aligned} X^T &= \{x \in X : (x, Tx) \in E(G)\} \\ X^S &= \{x \in X : (x, Sx) \in E(G)\} \end{aligned}$$

Remark 1.11. If the pair (T, S) is *G-orbital-cyclic pair*, then $X^T \neq \emptyset \iff X^S \neq \emptyset$.

Proof. Let $x_0 \in X^T$. Then $(x_0, Tx_0) \in E(G) \implies (Tx_0, STx_0) \in E(G)$.

If we denote by $x_1 = Tx_0$ we have that $(x_1, Sx_1) \in E(G)$, and thus, $X^S \neq \emptyset$. \square

2. Reich-Rus type operators

Theorem 2.1. Let T, S be two self-mappings on a complete extended b-metric space (X, d_θ) endowed with a directed graph G such that the pair (T, S) forms a *G-orbital-cyclic pair*. Suppose that

(i) $X^T \neq \emptyset$;

(ii) for all $x \in X^T$ and $y \in X^S$ and $k_1, k_2, k_3 > 0$, with $k_1 + k_2 + k_3 < 1$

$$d_\theta(Tx, Sy) \leq k_1 d_\theta(x, y) + k_2 d_\theta(x, Tx) + k_3 d_\theta(y, Sy);$$

(iii) for any sequence $(x_n)_{n \in \mathbb{N}} \subset X$, with $(x_n, x_{n+1}) \in E(G)$,

$$\lim_{n, m \rightarrow \infty} \theta(x_n, x_m) < \frac{1}{q}, \text{ where } q = \max \left\{ \frac{k_1 + k_2}{1 - k_3}, \frac{k_1 + k_3}{1 - k_2} \right\};$$

(iv) S and T are continuous,

or

(iv*) for any sequence $(x_n)_{n \in \mathbb{N}} \subset X$, with $x_n \rightarrow u$ as $n \rightarrow \infty$, and $(x_n, x_{n+1}) \in E(G)$, for $n \in \mathbb{N}$, we have, $u \in X^T \cap X^S$.

In these conditions $CFix(T, S) \neq \emptyset$.

Moreover, if we suppose

(v) if $(u, v) \in CFix(T, S)$ implies $u \in X^T$ and $v \in X^S$ then the pair (T, S) has a unique common fixed point.

Proof. Let $x_0 \in X^T$. Thus $(x_0, Tx_0) \in E(G)$.

Because the pair (T, S) is G -orbital-cyclic, we have $(Tx_0, STx_0) \in E(G)$.

If we denote by $x_1 = Tx_0$ we have $(x_1, Sx_1) \in E(G)$ and from here $(Sx_1, TSx_1) \in E(G)$. Denoting by $x_2 = Sx_1$ we have $(x_2, Tx_2) \in E(G)$.

By this procedure we construct inductively, a sequence $(x_n)_{n \in \mathbb{N}}$, with $x_{2n} = Sx_{2n-1}$ and $x_{2n+1} = Tx_{2n}$, such that $(x_{2n}, x_{2n+1}) \in E(G)$.

We shall suppose that $x_n \neq x_{n+1}$.

If, there exists $n_0 \in \mathbb{N}$, such that $x_{n_0} = x_{n_0+1}$, then, because $\Delta \subset E(G)$, $(x_{n_0}, x_{n_0+1}) \in E(G)$ and $u = x_{n_0}$ is a fixed point of T .

In order to show that $u \in CFix(T, S)$, we shall consider two cases for n_0 .

If $n_0 = 2k$, then $x_{2k} = x_{2k+1} = Tx_{2k}$ and thus, x_{2k} is a fixed point for T . Suppose that $d_\theta(Tx_{2k}, Sx_{2k+1}) > 0$, and let $x = x_{2k} \in X^T$ and $y = x_{2k+1} \in X^S$.

$$\begin{aligned} 0 &< d_\theta(x_{2k+1}, x_{2k+2}) = d_\theta(Tx_{2k}, Sx_{2k+1}) \leq k_1 d_\theta(x_{2k}, x_{2k+1}) + k_2 d_\theta(x_{2k}, Tx_{2k}) + k_3 d_\theta(x_{2k+1}, Sx_{2k+1}) \\ &= k_3 d_\theta(x_{2k+1}, Sx_{2k+1}) = k_3 d_\theta(x_{2k+1}, x_{2k+2}). \end{aligned}$$

In this way we reach to a contradiction.

In the same way we can prove the case $n_0 = 2k + 1$.

In conclusion $x_n \neq x_{n+1}$, for all $n \in \mathbb{N}$.

Now we shall prove that $(x_n)_{n \in \mathbb{N}}$ is a Cauchy sequence. In order to do this, we shall consider two possible cases:

Case 1. $x = x_{2n} \in X^T$ and $y = x_{2n+1} \in X^S$.

$$\begin{aligned} 0 &< d_\theta(x_{2n+1}, x_{2n+2}) = d_\theta(Tx_{2n}, Sx_{2n+1}) \\ &\leq k_1 d_\theta(x_{2n}, x_{2n+1}) + k_2 d_\theta(x_{2n}, Tx_{2n}) + k_3 d_\theta(x_{2n+1}, Sx_{2n+1}) \\ &= k_1 d_\theta(x_{2n}, x_{2n+1}) + k_2 d_\theta(x_{2n}, x_{2n+1}) + k_3 d_\theta(x_{2n+1}, x_{2n+2}) \end{aligned}$$

$$\begin{aligned} (1 - k_3) d_\theta(x_{2n+1}, x_{2n+2}) &\leq (k_1 + k_2) d_\theta(x_{2n}, x_{2n+1}) \\ d_\theta(x_{2n+1}, x_{2n+2}) &\leq \frac{k_1 + k_2}{1 - k_3} d_\theta(x_{2n}, x_{2n+1}) \\ d_\theta(x_{2n+1}, x_{2n+2}) &\leq q d_\theta(x_{2n}, x_{2n+1}). \end{aligned}$$

Case 2. $x = x_{2n} \in X^T$ and $y = x_{2n-1} \in X^S$.

$$\begin{aligned} 0 &< d_\theta(x_{2n+1}, x_{2n}) = d_\theta(Tx_{2n}, Sx_{2n-1}) \\ &\leq k_1 d_\theta(x_{2n}, x_{2n-1}) + k_2 d_\theta(x_{2n}, Tx_{2n}) + k_3 d_\theta(x_{2n-1}, Sx_{2n-1}) \\ &= k_1 d_\theta(x_{2n}, x_{2n-1}) + k_2 d_\theta(x_{2n}, x_{2n+1}) + k_3 d_\theta(x_{2n-1}, x_{2n}) \end{aligned}$$

$$\begin{aligned} (1 - k_2) d_\theta(x_{2n}, x_{2n+1}) &\leq (k_1 + k_3) d_\theta(x_{2n-1}, x_{2n}) \\ d_\theta(x_{2n}, x_{2n+1}) &\leq \frac{k_1 + k_3}{1 - k_2} d_\theta(x_{2n-1}, x_{2n}) \\ d_\theta(x_{2n}, x_{2n+1}) &\leq q d_\theta(x_{2n-1}, x_{2n}). \end{aligned}$$

In this way we have proved that

$$d_\theta(x_m, x_{m+1}) \leq q d_\theta(x_{m-1}, x_m), \text{ for all } m \in \mathbb{N}.$$

From Lemma 1.8., taking into account (iii), we obtain that $(x_m)_{m \in \mathbb{N}}$ is a Cauchy sequence in a complete extended b-metric space. Therefore, there is some point $u \in X$, such that $\lim_{m \rightarrow \infty} x_m = u$.

It is obvious that $\lim_{n \rightarrow \infty} x_{2n} = \lim_{n \rightarrow \infty} x_{2n+1} = u$.

Using (iv) we have

$$\begin{aligned} u &= \lim_{n \rightarrow \infty} x_{2n+1} = \lim_{n \rightarrow \infty} T(x_{2n}) = Tu, \\ u &= \lim_{n \rightarrow \infty} x_{2n+2} = \lim_{n \rightarrow \infty} S(x_{2n+1}) = Su. \end{aligned}$$

Hence $u \in CFix(T, S)$.

Let us suppose now that (iv*) take place, and let $x = u \in X^T$ and $y = x_{2n+1} \in X^S$.

$$\begin{aligned} 0 < d_\theta(Tu, x_{2n+2}) &= d_\theta(Tu, Sx_{2n+1}) \leq k_1 d_\theta(u, x_{2n+1}) + k_2 d_\theta(u, Tu) + k_3 d_\theta(x_{2n+1}, Sx_{2n+1}) \\ &= k_1 d_\theta(u, x_{2n+1}) + k_2 d_\theta(u, Tu) + k_3 d_\theta(x_{2n+1}, x_{2n+2}) = k_2 d_\theta(u, Tu). \end{aligned}$$

From here, we obtain that $d_\theta(u, Tu) = 0$.

Let now consider $x = x_{2n+1} \in X^T$ and $y = u \in X^S$.

$$\begin{aligned} 0 < d_\theta(x_{2n+1}, Su) &= d_\theta(Tx_{2n}, Su) \leq k_1 d_\theta(x_{2n}, u) + k_2 d_\theta(x_{2n}, Tx_{2n}) + k_3 d_\theta(u, Su) \\ &= k_1 d_\theta(u, x_{2n}) + k_2 d_\theta(x_{2n}, x_{2n+1}) + k_3 d_\theta(u, Su) = k_3 d_\theta(u, Su). \end{aligned}$$

From here, we obtain that $d_\theta(u, Su) = 0$, and thus, $u \in CFix(T, S)$.

Let us prove now the uniqueness of the common fixed point. Suppose that, there exist $u, v \in CFix(T, S)$, $u \neq v$. From (v) we have that $(u, Tu) \in E(G)$ and $(v, Sv) \in E(G)$. Now, using (ii) we obtain

$$0 < d_\theta(u, v) = d_\theta(Tu, Sv) \leq k_1 d_\theta(u, v) + k_2 d_\theta(u, Tu) + k_3 d_\theta(v, Sv) = k_1 d_\theta(u, v),$$

which is a contradiction. In conclusion $u = v$. □

3. Ciric type operators

Theorem 3.1. *Let T, S be two self-mappings on a complete extended b-metric space (X, d_θ) endowed with a directed graph G such that the pair (T, S) forms a G -orbital-cyclic pair. Suppose that*

(i) $X^T \neq \emptyset$;

(ii) for all $x \in X^T$ and $y \in X^S$ and $k_1, k_2, k_3 > 0$, with $k_1 + k_2 + k_3 < 1$

$$d_\theta(Tx, Sy) \leq k \max \{d_\theta(x, y), d_\theta(x, Tx), d_\theta(y, Sy)\};$$

(iii) for any sequence $(x_n)_{n \in \mathbb{N}} \subset X$, with $(x_n, x_{n+1}) \in E(G)$, $\lim_{n, m \rightarrow \infty} \theta(x_n, x_m) < \frac{1-k}{k}$;

(iv) S and T are continuous,

or

(iv*) for any sequence $(x_n)_{n \in \mathbb{N}} \subset X$, with $x_n \rightarrow u$ as $n \rightarrow \infty$, and $(x_n, x_{n+1}) \in E(G)$, for $n \in \mathbb{N}$, we have, $u \in X^T \cap X^S$.

In these conditions $CFix(T, S) \neq \emptyset$.

Moreover, if we suppose

(v) if $(u, v) \in CFix(T, S)$ implies $u \in X^T$ and $v \in X^S$ then the pair (T, S) has a unique common fixed point.

Proof. Let $x_0 \in X^T$. Just like in the proof of Theorem 2.1., we construct inductively, a sequence $(x_n)_{n \in \mathbb{N}}$, with $x_{2n} = Sx_{2n-1}$ and $x_{2n+1} = Tx_{2n}$, such that $(x_{2n}, x_{2n+1}) \in E(G)$.

We shall suppose that $x_n \neq x_{n+1}$.

If, there exists $n_0 \in \mathbb{N}$, such that $x_{n_0} = x_{n_0+1}$, then, because $\Delta \subset E(G)$, $(x_{n_0}, x_{n_0+1}) \in E(G)$ and $u = x_{n_0}$ is a fixed point of T .

In order to show that $u \in CFix(T, S)$, we shall consider two cases for n_0 .

If $n_0 = 2k$, then $x_{2k} = x_{2k+1} = Tx_{2k}$ an thus, x_{2k} is a fixed point for T . Suppose that $d_\theta(Tx_{2k}, Sx_{2k+1}) > 0$, and let $x = x_{2k} \in X^T$ and $y = x_{2k+1} \in X^S$.

$$\begin{aligned} 0 &< d_\theta(x_{2k+1}, x_{2k+2}) = d_\theta(Tx_{2k}, Sx_{2k+1}) \\ &\leq k \max \{d_\theta(x_{2k}, x_{2k+1}), d_\theta(x_{2k}, Tx_{2k}), d_\theta(x_{2k+1}, Sx_{2k+1})\} \\ &= kd_\theta(x_{2k+1}, Sx_{2k+1}) = kd_\theta(x_{2k+1}, x_{2k+2}). \end{aligned}$$

In this way we reach to a contradiction.

In the same way we can prove the case $n_0 = 2k + 1$.

In conclusion $x_n \neq x_{n+1}$, for all $n \in \mathbb{N}$.

Now we shall prove that $(x_n)_{n \in \mathbb{N}}$ is a Cauchy sequence. In order to do this, we shall consider two possible cases:

Case 1. $x = x_{2n} \in X^T$ and $y = x_{2n+1} \in X^S$.

$$\begin{aligned} 0 &< d_\theta(x_{2n+1}, x_{2n+2}) = d_\theta(Tx_{2n}, Sx_{2n+1}) \\ &\leq k \max \{d_\theta(x_{2n}, x_{2n+1}), d_\theta(x_{2n}, Tx_{2n}), d_\theta(x_{2n+1}, Sx_{2n+1})\} \\ &= k \max \{d_\theta(x_{2n}, x_{2n+1}), d_\theta(x_{2n+1}, x_{2n+2})\} \\ &\leq k [d_\theta(x_{2n}, x_{2n+1}) + d_\theta(x_{2n+1}, x_{2n+2})] \end{aligned}$$

$$\begin{aligned} (1-k) d_\theta(x_{2n+1}, x_{2n+2}) &\leq kd_\theta(x_{2n}, x_{2n+1}) \\ d_\theta(x_{2n+1}, x_{2n+2}) &\leq \frac{k}{1-k} d_\theta(x_{2n}, x_{2n+1}). \end{aligned}$$

Case 2. $x = x_{2n} \in X^T$ and $y = x_{2n-1} \in X^S$.

$$\begin{aligned} 0 &< d_\theta(x_{2n+1}, x_{2n}) = d_\theta(Tx_{2n}, Sx_{2n-1}) \\ &\leq k \max \{d_\theta(x_{2n}, x_{2n-1}), d_\theta(x_{2n}, Tx_{2n}), d_\theta(x_{2n-1}, Sx_{2n-1})\} \\ &= k \max \{d_\theta(x_{2n}, x_{2n-1}), d_\theta(x_{2n}, x_{2n+1})\} \\ &\leq k [d_\theta(x_{2n}, x_{2n-1}) + d_\theta(x_{2n}, x_{2n+1})] \end{aligned}$$

$$d_\theta(x_{2n}, x_{2n+1}) \leq \frac{k}{1-k} d_\theta(x_{2n-1}, x_{2n})$$

In this way we have proved that

$$d_\theta(x_m, x_{m+1}) \leq \frac{k}{1-k} d_\theta(x_{m-1}, x_m), \text{ for all } m \in \mathbb{N}.$$

From Lemma 1.8., taking into account (iii), we obtain that $(x_m)_{m \in \mathbb{N}}$ is a Cauchy sequence in a complete extended b-metric space. Therefore, there is some point $u \in X$, such that $\lim_{m \rightarrow \infty} x_m = u$.

It is obvious that $\lim_{n \rightarrow \infty} x_{2n} = \lim_{n \rightarrow \infty} x_{2n+1} = u$.

Using (iv) we have

$$\begin{aligned} u &= \lim_{n \rightarrow \infty} x_{2n+1} = \lim_{n \rightarrow \infty} T(x_{2n}) = Tu, \\ u &= \lim_{n \rightarrow \infty} x_{2n+2} = \lim_{n \rightarrow \infty} S(x_{2n+1}) = Su. \end{aligned}$$

Hence $u \in CFix(T, S)$.

Let us suppose now that (iv^*) take place, and let $x = u \in X^T$ and $y = x_{2n+1} \in X^S$.

$$0 < d_\theta(Tu, x_{2n+2}) = d_\theta(Tu, Sx_{2n+1}) \leq k \max \{d_\theta(u, x_{2n+1}), d_\theta(u, Tu), d_\theta(x_{2n+1}, Sx_{2n+1})\} = kd_\theta(u, Tu)$$

From here, we obtain that $d_\theta(u, Tu) = 0$.

Let now consider $x = x_{2n+1} \in X^T$ and $y = u \in X^S$.

$$0 < d_\theta(x_{2n+1}, Su) = d_\theta(Tx_{2n}, Su) \leq k \max \{d_\theta(x_{2n}, u), d_\theta(x_{2n}, Tx_{2n}), d_\theta(u, Su)\} = kd_\theta(u, Su)$$

From here, we obtain that $d_\theta(u, Su) = 0$, and thus, $u \in CFix(T, S)$.

Let us prove now the uniqueness of the common fixed point. Suppose that, there exist $u, v \in CFix(T, S)$, $u \neq v$. From (v) we have that $(u, Tu) \in E(G)$ and $(v, Sv) \in E(G)$. Now, using (ii) we obtain

$$0 < d_\theta(u, v) = d_\theta(Tu, Sv) \leq k \max \{d_\theta(u, v), d_\theta(u, Tu), d_\theta(v, Sv)\} = kd_\theta(u, v),$$

which is a contradiction. In conclusion $u = v$. □

References

- [1] B. Alqahtani, A. Fulga, E. Karapinar, Common fixed point results on extended b-metric space, *Journal of Inequalities and Applications*, (2018) 2018:158. <https://doi.org/10.1186/s13660-018-1745-4>
- [2] V. Berinde, Generalized contractions in quasimetric spaces, *Seminar on Fixed Point Theory*, Preprint no. 3(1993), 3-9.
- [3] S. Czerwik, Nonlinear set-valued contraction mappings in b-metric spaces, *Atti Sem. Mat. Univ. Modena*, 46(1998), 263-276.
- [4] T. Kamran, M. Samreen, O.U. Aim, A generalisation of b-metric space and some fixed point theorems, *Mathematics*, 5, 19(2017).