Results in Nonlinear Analysis **5** (2022) No. 1, 29–41.
https://doi.org/10.53006/rna.928654
Available online at www.resultsinnonlinearanalysis.com

Results in Nonlinear Analysis
ISSN 2636-7556

Peer Reviewed Scientific Journal

Sequential fractional pantograph differential equations with nonlocal boundary conditions: Uniqueness and Ulam-Hyers-Rassias stability

Mohamed Houas[a]

[a] *Laboratory FIMA, Khemis Miliana University, Khemis Miliana, Algeria.*

Abstract

In the current manuscript, we study the uniqueness and Ulam-stability of solutions for sequential fractional pantograph differential equations with nonlocal boundary conditions. The uniqueness of solutions is established by Banach's fixed point theorem. We also define and study the Ulam-Hyers stability and the Ulam-Hyers-Rassias stability of mentioned problem. An example is presented to illustrate the main results.

Keywords: Caputo derivative fixed point existence pantograph equations Ulam stability.
2010 MSC: 26A33 34A08 34B15.

1. Introduction

Differential equations of arbitrary order have recently been studied by many researchers, these equations will be used to describe phenomena of real world problems. For more details, see the works [2, 14, 21, 27] and the references therein. Many interesting and important area concerning of research for differential equations with fractional calculus are devoted to the existence theory and stability analysis of the solutions, for instance, for instance, see papers [4, 7, 9, 12, 17, 24]. Recently, several scholars have discussed the existence, uniqueness and different types of Ulam-stability of solutions for some classes of differential equations involving fractional derivatives, for instance, see [6, 13, 15, 16, 26] and the references cited therein. Considerable attention has been given to the study of the existence, uniqueness and Ulam stability of solutions for sequential fractional differential equations, we refer the reader to the monographs [20, 22, 29] and the reference therein. In the present work, we shall be concerned with a very special delay differential equation that has many applications

Email address: m.houas.st@univ-dbkm.dz (Mohamed Houas)

Received April 27, 2021; Accepted: January 8, 2022; Online: January 11, 2022

in different fields of pure and applied mathematics. The equation is called pantograph equation. For more information and some applications on pantograph equation, we refer the reader to the works [8, 10, 11, 23]. It is to note that the standard form of pantograph equation has the form

$$u^{'}(t) = Au(t) + Bu(\eta t),\ t \in [0,T],\ 0 < \eta < 1,$$

$$u(0) = u_0.$$

In recent years, many researchers have discussed the existence, uniqueness and different types of Ulam-Hyers stability of the above equation, For more details, see the monographs [1, 3, 18, 25, 28] and the references therein. In [5] the authors considered the following problem of the pantograph type

$${}^{C}D^{\alpha}u(t) = \phi(t, u(t), u(\eta t)), t \in [0,T],\ 0 < \alpha, \eta < 1,$$

$$u(0) = u_0,$$

where ${}^{C}D^{q}$ is the Caputo fractional derivative. The uniqueness results were obtained by applying Banach's contraction mapping principle. In this paper, we discuss the uniqueness, Ulam-Hyers stability and Ulam-Hyers-Rassias stability of solutions for the following sequential fractional pantograph equation

$$\left[D^{\alpha} + kD^{\beta}\right] u(t) = \phi\left(t, u(t), u(\eta t), D^{\beta}u(\eta t)\right), t \in [0,T], \tag{1}$$

under the conditions

$$u(0) = f(u), \quad u(T) = \theta,\ \theta \in \mathbb{R}, \tag{2}$$

where $k \in \mathbb{R}^{+}, 0 < \eta < 1, 1 < \alpha \leq 2, 0 < \beta \leq 1$, D^{α}, D^{β} are the Caputo type fractional derivatives, $\phi : [0,T] \times \mathbb{R} \times \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ and $f : C([0,T], \mathbb{R}) \to \mathbb{R}$ are given continuous functions. The operator D^{ϑ} is the fractional derivative in the sense of Caputo, defined by

$$\begin{aligned} D^{\vartheta}\varphi(t) &= \frac{1}{\Gamma(n-\vartheta)} \int_0^t (t-s)^{n-\vartheta-1} \varphi^{(n)}(s)\, ds \\ &= J^{n-\vartheta}\varphi^{(n)}(t), t > 0,\ n-1 < \vartheta < n,\ n \in N^{*}, \end{aligned}$$

and the Riemann-Liouville fractional integral of order $\vartheta > 0$, defined by

$$J^{\vartheta}\varphi(t) = \frac{1}{\Gamma(\vartheta)} \int_0^t (t-s)^{\vartheta-1} \varphi(s)\, ds, t > 0,$$

where $\Gamma(\vartheta) = \int_0^{\infty} e^{-u} x^{\vartheta-1} dx$.

Now, we give the following lemmas [19, 21]:

Lemma 1.1. *Let $m, s > 0$, $\varphi \in L_1([a,b])$. Then $J^m J^s \varphi(t) = J^{m+s}\varphi(t)$, $D^s J^s \varphi(t) = \varphi(t)$, $t \in [a,b]$.*

Lemma 1.2. *Let $s > m > 0$, $\varphi \in L_1([a,b])$. Then $D^m J^s \varphi(t) = J^{s-m}\varphi(t)$, $t \in [a,b]$.*

Also we give the following lemmas [19]:

Lemma 1.3. *For $\vartheta > 0$, the general solution of the fractional differential equation $D^{\vartheta}u(t) = 0$ is given by*

$$u(t) = \sum_{i=0}^{n-1} c_i t^i,$$

where $c_i \in \mathbb{R}$, $i = 0, 1, 2, .., n-1$, $n = [\vartheta] + 1$.

Lemma 1.4. *Let $\vartheta > 0$. Then*

$$J^{\vartheta} D^{\vartheta} u(t) = u(t) + \sum_{i=0}^{n-1} c_i t^i,$$

for some $c_i \in \mathbb{R}$, $i = 0, 1, 2, ..., n-1$, $n = [\vartheta] + 1$.

Let us now define the space $W = \left\{u : u \in C([0,T], \mathbb{R}), D^{\beta} u \in C([0,T], \mathbb{R})\right\}$ equipped, with the norm $\|u\|_W = 2\|u\| + \left\|D^{\beta} u\right\|$, where

$$\|u\| = \sup_{t \in [0,T]} |u(t)| \text{ and } \left\|D^{\beta} u\right\| = \sup_{t \in [0,T]} \left|D^{\beta} u(t)\right|.$$

It is clear that $(W, \|u\|_W)$ is a Banach space.

In that follows, we present the Ulam stability for the sequential fractional pantograph differential equations (1).

Definition 1.5. *The sequential fractional pantograph differential equations (1) is Ulam-Hyers stable if there exists a real number $\lambda_{\phi} > 0$ such that for each $\mu > 0$ and for each solution $v \in W$ of the inequality*

$$\left|\left[D^{\alpha} + kD^{\beta}\right] v(t) - \phi\left(t, v(t), v(\eta t), D^{\beta} v(\eta t)\right)\right| \leq \mu, \ t \in [0,T], \tag{3}$$

there exists a solution $u \in W$ of the sequential fractional pantograph differential equations (1) with

$$|v(t) - u(t)| \leq \lambda_{\phi} \mu, \ t \in [0,T].$$

Definition 1.6. *The sequential fractional pantograph differential equations (1) is generalized Ulam-Hyers stable if there exists $g_{\phi} \in C(\mathbb{R}_+, \mathbb{R}_+)$, $g_{\phi}(0) = 0$, such that for each solution $v \in W$ of the inequality (3), there exists a solution $u \in W$ of the sequential fractional pantograph differential equations (1) with*

$$|v(t) - u(t)| \leq g_{\phi}(\mu), \ t \in [0,T].$$

Definition 1.7. *The sequential fractional pantograph differential equations (1) is Ulam-Hyers-Rassias stable with respect to $h \in W$ if there exists a real number $\lambda_{\varphi} > 0$ such that for each $\mu > 0$ and for each solution $v \in W$ of the inequality*

$$\left|\left[D^{\alpha} + kD^{\beta}\right] v(t) - \phi\left(t, v(t), v(\eta t), D^{\beta} v(\eta t)\right)\right| \leq \mu h(t), \ t \in [0,T], \tag{4}$$

there exists a solution $u \in W$ of sequential fractional pantograph differential equations (1) with

$$|v(t) - u(t)| \leq \lambda_{\varphi} \mu h(t), \ t \in [0,T].$$

Definition 1.8. *The sequential fractional pantograph differential equations (1) is generalized Ulam-Hyers-Rassias stable with respect to $h \in C(J, \mathbb{R}_+)$ if there exists a real number $\lambda_{\phi,h} > 0$ such that for each solution $v \in W$ of the inequality*

$$\left|\left[D^{\alpha} + kD^{\beta}\right] v(t) - \phi\left(t, v(t), v(\eta t), D^{\beta} v(\eta t)\right)\right| \leq h(t), \ t \in [0,T], \tag{5}$$

there exists a solution $u \in W$ of the sequential fractional pantograph differential equations (1) with

$$|v(t) - u(t)| \leq \lambda_{\phi,h} h(t), \ t \in [0,T].$$

Remark 1.9. *A function $v \in W$ is a solution of the inequality (3) if and only if there exists a function $\psi : [0,T] \to \mathbb{R}$ (which depend on v) such that*

(1) : $|\psi(t)| \leq \mu, t \in [0,T]$.

(2) : $\left[D^{\alpha} + kD^{\beta}\right] v(t) = \phi\left(t, v(t), v(\eta t), D^{\beta} v(\eta t)\right) + \psi(t), t \in [0,T]$.

2. Main Results

We prove the following auxiliary lemma which is pivotal to define the solution for the problem (1)-(2).

Lemma 2.1. *Suppose that $h \in C([0,T],\ \mathbb{R})$ and consider the problem*

$$\begin{cases} \left[D^{\alpha}+kD^{\beta}\right]u(t)=h(t),\ t\in[0,T], \\ u(0)=f(u), \qquad u(T)=\theta,\theta\in\mathbb{R}, \\ k>0,\ 1<\alpha\leq 2,\ 0<\beta\leq 1. \end{cases} \tag{6}$$

Then, we have

$$\begin{aligned} u(t) &= \frac{1}{\Gamma(\alpha-\beta)}\int_0^t (t-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x (x-s)^{\beta-1}h(s)\,ds\right)dx \\ &\quad -\frac{k}{\Gamma(\alpha-\beta)}\int_0^t (t-x)^{\alpha-\beta-1}u(x)\,dx \\ &\quad +\frac{t^{\alpha-\beta}}{T^{\alpha-\beta}}\left[\theta-\frac{1}{\Gamma(\alpha-\beta)}\int_0^T (T-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x (x-s)^{\beta-1}h(s)\,ds\right)dx\right. \\ &\quad \left.+\frac{k}{\Gamma(\alpha-\beta)}\int_0^T (T-x)^{\alpha-\beta-1}u(x)\,dx\right]-\left[\frac{t^{\alpha-\beta}}{T^{\alpha-\beta}}-1\right]f(u). \end{aligned} \tag{7}$$

Proof. We have

$$\left[D^{\alpha}+kD^{\beta}\right]u(t)=h(t). \tag{8}$$

Now, writing the linear sequential fractional pantograph differential equation in (8) as

$$D^{\beta}\left[D^{\alpha-\beta}+k\right]u(t)=h(t). \tag{9}$$

By taking the Riemann-Liouville fractional integral of order β for (9), we get

$$\begin{aligned} u(t) &= \frac{1}{\Gamma(\alpha-\beta)}\int_0^t (t-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x (x-s)^{\beta-1}h(s)\,ds\right)dx \\ &\quad -\frac{k}{\Gamma(\alpha-\beta)}\int_0^t (t-x)^{\alpha-\beta-1}u(x)\,dx+\frac{t^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}c_0+d_0, \end{aligned} \tag{10}$$

where c_0 and d_0 are arbitrary constants. By the boundary condition $u(0)=f(u)$, we conclude that $d_0=f(u)$.

Using the boundary condition $u(T)=\theta$, we obtain that

$$\begin{aligned} c_0 &= \frac{\Gamma(\alpha-\beta+1)}{T^{\alpha-\beta}}\left[\theta-\frac{1}{\Gamma(\alpha-\beta)}\int_0^T (T-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x (x-s)^{\beta-1}h(s)\,ds\right)dx\right. \\ &\quad \left.+\frac{k}{\Gamma(\alpha-\beta)}\int_0^T (T-x)^{\alpha-\beta-1}u(x)\,dx-f(u)\right]. \end{aligned} \tag{11}$$

Substituting the values of c_0 and c_1 in (10), we obtain the solution (7). This completes the proof.

□

In view of Lemma 2.1, we define the operator $O : W \to W$ by:

$$\begin{aligned} Ou(t) &= \frac{1}{\Gamma(\alpha-\beta)}\int_0^t (t-x)^{\alpha-\beta-1} \\ &\times\left(\frac{1}{\Gamma(\beta)}\int_0^x (x-s)^{\beta-1}\phi\left(t,u(s),u(\eta s),D^{\beta}u(\eta s)\right)ds\right)dx \\ &-\frac{k}{\Gamma(\alpha-\beta)}\int_0^t (t-x)^{\alpha-\beta-1}u(x)\,dx \\ &+\frac{t^{\alpha-\beta}}{T^{\alpha-\beta}}\left[\theta-\frac{1}{\Gamma(\alpha-\beta)}\int_0^T (T-x)^{\alpha-\beta-1}\right. \\ &\times\left(\frac{1}{\Gamma(\beta)}\int_0^x (x-s)^{\beta-1}\phi\left(s,u(s),u(\eta s),D^{\beta}u(\eta s)\right)ds\right)dx \\ &\left.+\frac{k}{\Gamma(\alpha-\beta)}\int_0^T (T-x)^{\alpha-\beta-1}u(x)\,dx\right]-\left[\frac{t^{\alpha-\beta}}{T^{\alpha-\beta}}-1\right]f(u). \end{aligned} \tag{12}$$

In the sequel, we need the following hypotheses:

(H_1) : $\phi : J\times\mathbb{R}\times\mathbb{R}\times\mathbb{R}\to\mathbb{R}$ is continuous function and there exists nonnegative constant ω such that for all $t\in[0,1]$ and $y_i,\ z_i\in\mathbb{R}\ (i=1,2,3)$,

$$|\phi(t,y_1,y_2,y_3)-\phi(t,z_1,z_2,z_3)|\le\omega\left(|t_1-z_1|+|y_2-z_2|+|y_3-z_3|\right).$$

(H_2) : $f : C([0,T],\mathbb{R})\to\mathbb{R}$ is a continuous function with $f(0)=0$ and there exists constant $\varpi>0$ such that

$$|f(u)-f(v)|\le\varpi\|u-v\|,\ \forall u,v\in C([0,T],\mathbb{R}).$$

For convenience, we define:

$$\begin{aligned} \nabla_0 &: = \frac{\omega T^{\alpha}}{\Gamma(\alpha+1)}+\frac{kT^{\alpha}}{\Gamma(\alpha-\beta+1)}+\varpi, \\ \nabla_1 &: = \frac{\omega T^{\alpha-1}}{\Gamma(\alpha)}+\frac{(\alpha-\beta)\omega T^{\alpha-1}}{\Gamma(\alpha+1)}+\frac{kT^{\alpha-\beta-1}}{\Gamma(\alpha-\beta)}+\frac{(\alpha-\beta)kT^{\alpha-\beta-1}}{\Gamma(\alpha-\beta+1)}+\frac{\varpi(\alpha-\beta)}{T^{\alpha}}, \\ \Pi_0 &: = \frac{2T^{\alpha}L}{\Gamma(\alpha+1)}+|\theta|,\ \Pi_1=\frac{(\alpha-\beta)}{T^{\alpha}}|\theta|. \end{aligned} \tag{13}$$

2.1. Existence and uniqueness of solution

The result is concerned with the existence and uniqueness of the solution for the problem (1)-(2 and is based on Banach's fixed point theorem.

Theorem 2.2. *Assume that* (H_1) *and* (H_2) *hold. If the inequality*

$$\nabla_1 T^{1-\beta}<\Gamma(2-\beta)(1-4\nabla_0), \tag{14}$$

is valid, then the problem (1)-(2) has a unique solution on $[0,T]$.

Proof. Let us fix $\sup_{t\in[0,T]}\phi(t,0,0,0)=L<\infty$ and define

$$r\ge\frac{2\Pi_0+\frac{2T^{1-\beta}}{\Gamma(2-\beta)}\Pi_1}{1-4\nabla_0-\frac{T^{1-\beta}}{\Gamma(2-\beta)}\nabla_1}.$$

We show that $OB_r\subset B_r$, where O defined by (10) and $B_r=\{u\in W:\|u\|_W\le r\}$.

For $u \in B_r$, we have

$$\begin{aligned} & \left|\phi\left(t, u(t), u(\eta t), D^{\beta} u(\eta t)\right)\right| \\ \leq & \left(\left|\phi\left(t, u(t), u(\eta t), D^{\beta} u(\eta t)\right)-\phi(t, 0, 0, 0)\right|+|\varphi(t, 0, 0, 0)|\right) \\ & \left(\left|\phi\left(t, u(t), u(\eta t), D^{\beta} u(\eta t)\right)-\phi(t, 0, 0, 0)\right|\right)+|\varphi(t, 0, 0, 0)| \\ \leq & \omega\left(|u(t)|+|u(\eta t)|+\left|D^{\beta} u(t)\right|\right)+L \leq \omega\left(2\|u\|+\left\|D^{\beta} u\right\|\right)+L \\ \leq & \omega\|u\|_W+N \leq \omega r+L, \end{aligned} \tag{15}$$

and

$$|f(u)| \leq \varpi\|u\| \leq \varpi\|u\|_W \leq \varpi r. \tag{16}$$

It follows from (15) and (16), that

$$\begin{aligned} |O u(t)| \leq & \frac{1}{\Gamma(\alpha-\beta)} \int_0^t (t-x)^{\alpha-\beta-1} \\ & \times\left(\frac{1}{\Gamma(\beta)} \int_0^x (x-s)^{\beta-1}\left|\phi\left(t, u(s), u(\eta s), D^{\beta} u(\eta s)\right)\right| d s\right) d x \\ & +\frac{k}{\Gamma(\alpha-\beta)} \int_0^t (t-x)^{\alpha-\beta-1}|u(x)| d x \\ & +\frac{t^{\alpha-\beta}}{T^{\alpha-\beta}}\left[|\theta|+\frac{1}{\Gamma(\alpha-\beta)} \int_0^T (T-x)^{\alpha-\beta-1}\right. \\ & \times\left(\frac{1}{\Gamma(\beta)} \int_0^x (x-s)^{\beta-1}\left|\phi\left(s, u(s), u(\eta s), D^{\beta} u(\eta s)\right)\right| d s\right) d x \\ & \left.+\frac{k}{\Gamma(\alpha-\beta)} \int_0^T (T-x)^{\alpha-\beta-1}|u(x)| d x\right]+\left|\frac{t^{\alpha-\beta}}{T^{\alpha-\beta}}-1\right||f(u)| \\ \leq & 2\left(\frac{\omega T^{\alpha}}{\Gamma(\alpha+1)}+\frac{k T^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}+\varpi\right) r+2\left(\frac{T^{\alpha} L}{\Gamma(\alpha+1)}\right)+|\theta|. \\ = & 2 \nabla_0 r+\Pi_0. \end{aligned} \tag{17}$$

On the other hand, we have

$$\begin{aligned}
&\left|O\left(u\right)^{'}\left(t\right)\right| \leq \frac{1}{\Gamma\left(\alpha-\beta-1\right)}\int_0^t\left(t-x\right)^{\alpha-\beta-2} \\
&\left(\frac{1}{\Gamma\left(\beta\right)}\int_0^x\left(x-s\right)^{\beta-1}\left|\phi\left(t,u\left(s\right),u\left(\eta s\right),D^{\beta}u\left(\eta s\right)\right)\right|ds\right)dx \\
&+\frac{k}{\Gamma\left(\alpha-\beta-1\right)}\int_0^t\left(t-x\right)^{\alpha-\beta-2}\left|u\left(x\right)\right|dx \\
&+\frac{\left(\alpha-\beta\right)t^{\alpha-\beta-1}}{T^{\alpha-\beta}}\left[\left|\theta\right|+\frac{1}{\Gamma\left(\alpha-\beta\right)}\int_0^T\left(T-x\right)^{\alpha-\beta-1}\right. \\
&\left(\frac{1}{\Gamma\left(\beta\right)}\int_0^x\left(x-s\right)^{\beta-1}\left|\phi\left(s,u\left(s\right),u\left(\eta s\right),D^{\beta}u\left(\eta s\right)\right)\right|ds\right)dx \\
&\left.+\frac{k}{\Gamma\left(\alpha-\beta\right)}\int_0^T\left(T-x\right)^{\alpha-\beta-1}\left|u\left(x\right)\right|dx\right]+\frac{\left(\alpha-\beta\right)t^{\alpha-\beta-1}}{T^{\alpha-\beta}}\left|f\left(u\right)\right| \\
\leq\ &\left(\frac{\omega T^{\alpha-1}}{\Gamma\left(\alpha\right)}+\frac{\left(\alpha-\beta\right)\omega T^{\alpha}}{\Gamma\left(\alpha+1\right)}+\frac{kT^{\alpha-\beta-1}}{\Gamma\left(\alpha-\beta\right)}+\frac{\left(\alpha-\beta\right)kT^{\alpha-\beta-1}}{\Gamma\left(\alpha-\beta+1\right)}\right. \\
&\left.+\frac{\varpi\left(\alpha-\beta\right)}{T^{\alpha}}\right)r+\frac{\left(\alpha-\beta\right)}{T^{\alpha}}\left|\theta\right| \\
=\ &\nabla_1 r+\Pi_1,
\end{aligned} \tag{18}$$

which implies that

$$\left|D^{\beta}Ou\left(t\right)\right| \leq \int_0^t\frac{\left(t-s\right)^{-\beta}}{\Gamma\left(1-\beta\right)}\left|O\left(u\right)^{'}\left(t\right)\right|ds \leq \frac{T^{1-\beta}}{\Gamma\left(2-\beta\right)}\left(\nabla_1 r+\Pi_1\right). \tag{19}$$

Thus

$$\begin{aligned}
\left\|Ou\right\|_W &= 2\left\|Ou\right\|+\left\|D^{\beta}Ou\right\| \\
&\leq \left(4\nabla_0+\frac{T^{1-\beta}}{\Gamma\left(2-\beta\right)}\nabla_1\right)r+2\Pi_0+\frac{T^{1-\beta}}{\Gamma\left(2-\beta\right)}\Pi_1 \leq r,
\end{aligned} \tag{20}$$

which implies that $OB_r \subset B_r$. Now for $u,v \in B_r$ and for all $t \in [0,T]$, we obtain

$$\begin{aligned}
&\left|Ou\left(t\right)-Ov\left(t\right)\right| \\
\leq\ &\frac{1}{\Gamma\left(\alpha-\beta\right)}\int_0^t\left(t-x\right)^{\alpha-\beta-1}\left(\frac{1}{\Gamma\left(\beta\right)}\int_0^x\left(x-s\right)^{\beta-1}\right. \\
&\left.\left|\phi\left(s,u\left(s\right),u\left(\eta s\right),D^{\beta}u\left(\eta s\right)\right)-\phi\left(s,v\left(s\right),v\left(\eta s\right),D^{\beta}v\left(\eta s\right)\right)\right|ds\right)dx \\
&+\frac{1}{\Gamma\left(\alpha-\beta\right)}\int_0^t\left(t-x\right)^{\alpha-\beta-1}\left|u\left(x\right)-v\left(x\right)\right|dx \\
&+\frac{t^{\alpha-\beta}}{T^{\alpha-\beta}\Gamma\left(\alpha-\beta\right)}\int_0^T\left(T-x\right)^{\alpha-\beta-1}\left(\frac{1}{\Gamma\left(\beta\right)}\int_0^x\left(x-s\right)^{\beta-1}\right. \\
&\left.\left|\phi\left(s,u\left(s\right),u\left(\eta s\right),D^{\beta}u\left(\eta s\right)\right)-\phi\left(s,v\left(s\right),v\left(\eta s\right),D^{\beta}v\left(\eta s\right)\right)\right|ds\right)dx \\
&+\frac{t^{\alpha-\beta}}{T^{\alpha-\beta}\Gamma\left(\alpha-\beta\right)}\int_0^T\left(T-x\right)^{\alpha-\beta-1}\left|u\left(x\right)-v\left(x\right)\right|dx+\left|\frac{t^{\alpha-\beta}}{T^{\alpha-\beta}}-1\right|\left|f\left(u\right)-f\left(v\right)\right| \\
\leq\ &2\left[\frac{T^{\alpha}\omega}{\Gamma\left(\alpha+1\right)}+\frac{T^{\alpha-\beta}}{\Gamma\left(\alpha-\beta+1\right)}+\varpi\right]\left\|u-v\right\|_W \\
=\ &2\nabla_0\left\|u-v\right\|_W.
\end{aligned} \tag{21}$$

We also have

$$\begin{aligned}
&\left|O\left(u\right)^{'}\left(t\right)-O\left(v\right)^{'}\left(t\right)\right| \qquad (22)\\
\leq\ & \frac{1}{\Gamma\left(\alpha-\beta-1\right)}\int_0^t\left(t-x\right)^{\alpha-\beta-2}\left(\frac{1}{\Gamma\left(\beta\right)}\int_0^x\left(x-s\right)^{\beta-1}\right.\\
&\left.\left|\phi\left(s,u\left(s\right),u\left(\eta s\right),D^{\beta}u\left(\eta s\right)\right)-\phi\left(s,v\left(s\right),v\left(\eta s\right),D^{\beta}v\left(\eta s\right)\right)\right|ds\right)dx\\
&+\frac{1}{\Gamma\left(\alpha-\beta-1\right)}\int_0^t\left(t-x\right)^{\alpha-\beta-2}\left|u\left(x\right)-v\left(x\right)\right|dx\\
&+\frac{\left(\alpha-\beta\right)t^{\alpha-\beta-1}}{T^{\alpha-\beta}\Gamma\left(\alpha-\beta\right)}\int_0^T\left(T-x\right)^{\alpha-\beta-1}\left(\frac{1}{\Gamma\left(\beta\right)}\int_0^x\left(x-s\right)^{\beta-1}\right.\\
&\left.\left|\phi\left(s,u\left(s\right),u\left(\eta s\right),D^{\beta}u\left(\eta s\right)\right)-\phi\left(s,v\left(s\right),v\left(\eta s\right),D^{\beta}v\left(\eta s\right)\right)\right|ds\right)dx\\
&\frac{\left(\alpha-\beta\right)t^{\alpha-\beta-1}}{T^{\alpha-\beta}\Gamma\left(\alpha-\beta\right)}\int_0^T\left(T-x\right)^{\alpha-\beta-1}\left|u\left(x\right)-v\left(x\right)\right|dx+\frac{\left(\alpha-\beta\right)t^{\alpha-\beta}}{T^{\alpha-\beta}}\left|f\left(u\right)-f\left(v\right)\right|\\
\leq\ & \left(\frac{\omega T^{\alpha-1}}{\Gamma\left(\alpha\right)}+\frac{\left(\alpha-\beta\right)\omega T^{\alpha-1}}{\Gamma\left(\alpha+1\right)}+\frac{T^{\alpha-\beta-1}}{\Gamma\left(\alpha-\beta\right)}+\frac{\left(\alpha-\beta\right)T^{\alpha-\beta-1}}{\Gamma\left(\alpha-\beta+1\right)}\right.\\
&\left.+\frac{\varpi\left(\alpha-\beta\right)}{T^{\alpha}}\right)\left\|u-v\right\|_W\\
=\ & \nabla_1\left\|u-v\right\|_W.
\end{aligned}$$

Thus we obtain

$$\begin{aligned}
\left|D^{\beta}Ou\left(t\right)-D^{\beta}Ov\left(t\right)\right| &\leq \int_0^t\frac{\left(t-s\right)^{1-(\alpha+\beta)}}{\Gamma\left(2-\alpha-\beta\right)}\left|O\left(u\right)^{'}\left(t\right)-O\left(v\right)^{'}\left(t\right)\right|ds\\
&\leq \frac{\nabla_1 T^{1-\beta}}{\Gamma\left(2-\beta\right)}\left\|u-v\right\|_W. \qquad (23)
\end{aligned}$$

From the above inequalities, we get

$$\begin{aligned}
\left\|Ou-Ov\right\|_W &= 2\left\|Ou-Ov\right\|+\left\|D^{\beta}Ou-D^{\beta}Ov\right\|\\
&\leq \left(4\nabla_0+\frac{\nabla_1 T^{1-\beta}}{\Gamma\left(2-\beta\right)}\right)\left\|u-v\right\|_W. \qquad (24)
\end{aligned}$$

By (14), we see that O is a contractive operator. Consequently, by the Banach fixed point theorem, O has a fixed point which is a solution of system (1)-(2). This completes the proof. □

2.2. Ulam-Hyers-Rassias stability

In the following section, we will study Ulam's type stability of the sequential fractional pantograph differential equations (1).

Theorem 2.3. *Assume that $\phi:[0,T]\times\mathbb{R}\times\mathbb{R}\times\mathbb{R}\to\mathbb{R}$ is a continuous function satisfying (H_1). If*

$$\frac{kT^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}<1-\frac{\omega T^{\alpha}}{\Gamma\left(\alpha+1\right)}. \qquad (25)$$

Then the sequential fractional pantograph differential equations (1) is Ulam-Hyers stable and consequently, generalized Ulam-Hyers stable.

Proof. Let $v\in W$ be a solution of the inequality (3), i.e.

$$\left|\left[D^{\alpha}+kD^{\beta}\right]v\left(t\right)-\phi\left(t,v\left(t\right),v\left(\eta t\right),D^{\beta}v\left(\eta t\right)\right)\right|\leq\mu, t\in\left[0,T\right],$$

and let us denote by $u \in W$ the unique solution of the problem

$$\begin{aligned}\left[D^{\alpha}+kD^{\beta}\right]u(t) &= \phi\left(t,u(t),u(\eta t),D^{\beta}u(\eta t)\right),k>0,t\in[0,T],0<\eta<1,\\ u(0) &= v(0),\ u(T)=v(T).\end{aligned}$$

Thanks to Lemma 2.1, we can write

$$u(t)=I^{\alpha-\beta}\left[I^{\beta}h(t)\right]-kI^{\alpha-\beta}u(t)+c_0\frac{t^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}+d_0.$$

By integration of the inequality (3), we obtain

$$\begin{aligned}&\left|v(t)-I^{\alpha-\beta}\left[I^{\beta}h_v(t)\right]-kI^{\alpha-\beta}v(t)-c_1\frac{t^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}-d_1\right|\\ \leq\ &\frac{\mu t^{\alpha}}{\Gamma(\alpha+1)}\leq\frac{\mu T^{\alpha}}{\Gamma(\alpha+1)},\end{aligned}\tag{26}$$

where

$$h_v(t)=\phi\left(t,v(t),v(\eta t),D^{\beta}v(\eta t)\right).\tag{27}$$

On the other hand, if $u(0)=v(0)$ and $u(T)=v(T)$, then

$$c_0=c_1 \text{ and } d_0=d_1.$$

For any $t\in[0,T]$, we have

$$\begin{aligned}v(t)-u(t) &= v(t)-I^{\alpha-\beta}\left[I^{\beta}h_u(t)\right]-kI^{\alpha-\beta}v(t)-c_1\frac{t^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}-d_1\\ &\quad+I^{\alpha-\beta}\left[I^{\beta}\left(h_v(t)-h_u(t)\right)\right]-kI^{\alpha-\beta}\left(v(t)-u(t)\right),\end{aligned}\tag{28}$$

where

$$h_u(t)=\phi\left(t,u(t),u(\eta t),D^{\beta}u(\eta t)\right),\tag{29}$$

then

$$\begin{aligned}&I^{\alpha-\beta}\left[I^{\beta}\left(h_v(t)-h_u(t)\right)\right]\\ &=I^{\alpha-\beta}\left[I^{\beta}\left(\phi\left(t,u(t),u(\eta t),D^{\beta}u(\eta t)\right)-\phi\left(t,v(t),v(\eta t),D^{\beta}v(\eta t)\right)\right)\right]\\ &=\frac{1}{\Gamma(\alpha-\beta)}\int_0^t(t-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x(x-s)^{\beta-1}\phi\left(t,u(t),u(\eta t),D^{\beta}u(\eta t)\right)\right.\\ &\left.-\phi\left(t,v(t),v(\eta t),D^{\beta}v(\eta t)\right)ds\right)dx.\end{aligned}\tag{30}$$

By (H_1), we can write

$$\begin{aligned}&I^{\alpha-\beta}\left[I^{\beta}\left(h_v(t)-h_u(t)\right)\right]\\ \leq\ &\frac{1}{\Gamma(\alpha-\beta)}\int_0^t(t-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x(x-s)^{\beta-1}\omega\left\|v(s)-u(s)\right\|_W ds\right)dx.\end{aligned}\tag{31}$$

Using (31), we get

$$\begin{aligned}|v(t)-u(t)| &\leq \left|v(t)-I^{\alpha-\beta}\left[I^{\beta}h_u(t)\right]-kI^{\alpha-\beta}v(t)-c_1\frac{t^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}-d_1\right| \\ &+\frac{\omega}{\Gamma(\alpha-\beta)}\int_0^t(t-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x(x-s)^{\beta-1}\|v(s)-u(s)\|_W\,ds\right)dx \\ &+\frac{k}{\Gamma(\alpha-\beta)}\int_0^t(t-s)^{\alpha-\beta-1}\|v(s)-u(s)\|_W\,ds. \end{aligned} \tag{32}$$

Hence,

$$|v(t)-u(t)|\leq\frac{\mu T^{\alpha}}{\Gamma(\alpha+1)}+\left(\frac{\omega T^{\alpha}}{\Gamma(\alpha+1)}+\frac{kT^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}\right)\|v(s)-u(s)\|_W. \tag{33}$$

Then

$$\|v(s)-u(s)\|_W\left[1-\left(\frac{\omega T^{\alpha}}{\Gamma(\alpha+1)}+\frac{kT^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}\right)\right]\leq\frac{\mu T^{\alpha}}{\Gamma(\alpha+1)}. \tag{34}$$

For each $t\in[0,T]$

$$\|u(t)-v(t)\|\leq\frac{T^{\alpha}}{\Gamma(\alpha+1)\left[1-\left(\frac{\omega T^{\alpha}}{\Gamma(\alpha+1)}+\frac{kT^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}\right)\right]}\mu:=\lambda_{\phi}\mu. \tag{35}$$

Therefore, the sequential fractional pantograph differential equations (1) is Ulam-Hyers stable. By taking $g_{\phi}(\sigma)=\lambda_{\phi}\mu, g_{\phi}(0)=0$ yields that the sequential fractional pantograph differential equations (1) is generalized Ulam-Hyers stable. □

Now, we will study Ulam-Hyers-Rassias stability of the sequential fractional pantograph differential equations (1).

Theorem 2.4. *Let $\varphi:[0,T]\times\mathbb{R}\times\mathbb{R}\times\mathbb{R}\to\mathbb{R}$ be a continuous function and suppose that (H_1) and (25) hold. In addition, the following hypothesis holds*

(H_3) : *There exists an function $h\in C([0,T],\mathbb{R}_+)$ and there exists $\pi_h>0$ such that for any $t\in[0,T]$*

$$\frac{1}{\Gamma(\alpha-\beta)}\int_0^t(t-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x(x-s)^{\beta-1}h(s)\,ds\right)dx\leq\pi_h h(t).$$

Then the sequential fractional pantograph differential equations (1) is Ulam-Hyers-Rassias stable.

Proof. Let us denote by $v\in W$ the solution of the inequality (4), i.e.

$$\left|\left[D^{\alpha}+kD^{\beta}\right]v(t)-\phi\left(t,v(t),v(\eta t),D^{\beta}v(\eta t)\right)\right|\leq\mu h(t),\ t\in[0,T].$$

Let $u\in W$ be a the unique solution of the problem

$$\begin{aligned}\left[D^{\alpha}+kD^{\beta}\right]u(t) &= \phi\left(t,u(t),u(\eta t),D^{\beta}u(\eta t)\right),k>0,t\in[0,T],0<\eta<1, \\ u(0) &= v(0),\ u(T)=v(T). \end{aligned}$$

Applying Lemma 2.1, we get

$$u(t)=I^{\alpha-\beta}\left[I^{\beta}h(t)\right]-kI^{\alpha-\beta}u(t)+c_0\frac{t^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}+d_0.$$

Integrating the inequality (4), we can state that

$$\begin{aligned} &\left|v(t)-I^{\alpha-\beta}\left[I^{\beta}h_v(t)\right]-kI^{\alpha-\beta}v(t)-c_1\frac{t^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}-d_1\right| \\ \leq\ &\frac{\mu}{\Gamma(\alpha-\beta)}\int_0^t(t-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x(x-s)^{\beta-1}h(s)\,ds\right)dx, \end{aligned} \tag{36}$$

where $h_v(t)$ is given by (27).

So, by (H_1), we obtain

$$\begin{aligned} |v(t)-u(t)| \leq\ &\frac{\mu}{\Gamma(\alpha-\beta)}\int_0^t(t-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x(x-s)^{\beta-1}h(s)\,ds\right)dx \\ &+\frac{1}{\Gamma(\alpha-\beta)}\int_0^t(t-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x(x-s)^{\beta-1}\|v(s)-u(s)\|_W\,ds\right)dx \\ &+\frac{k}{\Gamma(\alpha-\beta)}\int_0^t(t-s)^{\alpha-\beta-1}\|v(s)-u(s)\|_W\,ds. \end{aligned} \tag{37}$$

Now, using (H_3), we have

$$|v(t)-u(t)|\leq\mu\pi_h h(t)+\left(\frac{\omega T^{\alpha}}{\Gamma(\alpha+1)}+\frac{kT^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}\right)\|v(s)-u(s)\|_W, \tag{38}$$

which implies that

$$\|v(s)-u(s)\|\left[1-\frac{\omega T^{\alpha}}{\Gamma(\alpha+1)}-\frac{kT^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}\right]\leq\mu\pi_h h(t). \tag{39}$$

For any $t\in[0,T]$, we have

$$\|u(t)-v(t)\|\leq\left[\frac{\pi_h}{1-\frac{\omega T^{\alpha}}{\Gamma(\alpha+1)}-\frac{kT^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}}\right]\mu h(t):=\lambda_{\phi}\mu h(t). \tag{40}$$

Then, the fractional boundary value problem (1) is Ulam-Hyers-Rassias stable. □

3. An example

To illustrate our main results, we treat the following example.

Example 3.1. *Let us consider the following fractional boundary value problem*

$$\begin{cases} \left(D^{\frac{5}{3}}+\frac{1}{10}D^{\frac{3}{4}}\right)u(t)=\frac{1}{2.20^2\pi}\sin(2\pi u(t))+\frac{1}{20^2}\sin(t)\,u\left(\frac{1}{2}t\right) \\ \qquad\qquad +\frac{1}{20^2}D^{\frac{3}{4}}u\left(\frac{1}{2}t\right)+\frac{5}{23},\ t\in[0,1], \\ x(0)=\frac{1}{60}\tan^{-1}u(t),\ u(1)=\frac{7}{5}. \end{cases} \tag{41}$$

For this example, we have

$$\begin{aligned} \varphi(t,u,v,w) &= \frac{1}{2.20^2\pi}\sin(2\pi u)+\frac{1}{20^2}\sin(t)\,u \\ +\frac{1}{20^2}w+\frac{5}{23},\ t &\in [0,1], \end{aligned}$$

and

$$f(u)=\frac{1}{60}\tan^{-1}u.$$

So, for any $(u_1,u_2,u_3),(v_1,v_2,v_3)\in\mathbb{R}^3$ *and* $t\in[0,1]$, *we can write*

$$|\varphi(t,u_1,v_1,w_1)-\varphi(t,u_2,v_2,w_2)|\leq\frac{1}{20^2}\left(|u_1-v_1|+|u_2-v_2|+|u_3-v_3|\right).$$

Hence the condition (H_1) *holds with* $\omega=\frac{1}{20^2}$. *Also for all* $u,v\in C([0,1],\mathbb{R})$, *we have*

$$|f(u)-f(v)|\leq\frac{1}{60}|u-v|.$$

So, (H_2) *is satisfied with* $\varpi=\frac{1}{60}$.

For $\alpha=\frac{5}{3},\beta=\frac{3}{4},k=\frac{1}{10},T=1$ *and* $\eta=\frac{1}{2}$, *we have*

$$\nabla_0=0.121\,68,\quad\nabla_1=0.209\,05.$$

Using the given data, we find that

$$\frac{\nabla_1T^{1-\beta}}{\Gamma(2-\beta)}=0.230\,64<1-4\nabla_0=0.513\,2.$$

Hence by Theorem 11, the problem (41) has a unique solution.

Also, we have

$$\frac{kT^{\alpha-\beta}}{\Gamma(\alpha-\beta+1)}=0.103\,35<1-\frac{\omega T^{\alpha}}{\Gamma(\alpha+1)}=0.998\,34.$$

Then, all the hypotheses of Theorem 12 are satisfied. Thus, by the conclusion of Theorem 12, problem (41) is Ulam-Hyers stable.

Let $h(t)=\gamma t^2,\gamma\in\mathbb{R}$. *We have*

$$\frac{1}{\Gamma(\alpha-\beta)}\int_0^t(t-x)^{\alpha-\beta-1}\left(\frac{1}{\Gamma(\beta)}\int_0^x(x-s)^{\beta-1}h(s)\,ds\right)dx\leq\frac{2\gamma}{\Gamma\left(\frac{14}{3}\right)}t^2=\pi_hh(t).$$

Thus condition (H_3) *is satisfied with* $h(t)=\gamma t^2$ *and* $\pi_h=\frac{2\gamma}{\Gamma\left(\frac{14}{3}\right)}$. *It follows from Theorem 13 problem (41) is Ulam-Hyers-Rassias stable.*

4. Conclusion

In this work, we have discussed the uniqueness and different types of Ulam-stability of solutions for sequential fractional pantograph differential equations with nonlocal boundary conditions. We have establish the uniqueness results applying the Banach contraction principle. Also, we have proved different types of Ulam stability results including Ulam-Hyers stability, generalized Ulam-Hyers stability and Ulam-Hyers-Rassias. For justification, a numerical example has been given to illustrate our main results.

References

[1] M.S. Abdo, T. Abdeljawad, K.D. Kucche, M.A. Alqudah, S.M. Ali and M.B. Jeelani, On nonlinear pantograph fractional differential equations with Atangana-Baleanu-Caputo derivative, Adv. Difference . Equ. 2021: 65 (2021), 1-17.

[2] M. S. Abdo, T. Abdeljawad, K. Shah and S. M. Ali, On nonlinear coupled evolution system with nonlocal subsidiary conditions under fractal-fractional order derivative, Math Meth Appl Sci. 44(8) (2021), 6581-6600.

[3] A. Ali, I. Mahariq, K. Shah, T. Abdeljawad and B. Al-Sheikh, Stability analysis of initial value problem of pantograph-type implicit fractional differential equations with impulsive conditions, Adv. Difference . Equ. 2021: 55 (2021), 1-17.

[4] B. Azizollah, Q.M. Al-Mdallal, On the existence of positive solutions for a non-autonomous fractional differential equation with integral boundary conditions, CMDE. 9(1) (2021), 36-51.

[5] K. Balachandran, S. Kiruthika and J.J. Trujillo, Existence of solutions of Nonlinear fractional pantograph equations, Acta Mathematica Scientia. 33B (2013), 1-9.

[6] W. Benhamida, S. Hamani and J. Henderson, Boundary value problems for Caputo-Hadamard fractional differential equations, Adv.Theory Nonlinear Anal. Appl. 2(3) (2018), 138-145
[7] A. Boutiara, M. S. Abdo, M. A. Alqudah and T. Abdeljawad, On a class of Langevin equations in the frame of Caputo function-dependent- kernel fractional derivatives with antiperiodic boundary conditions, AIMS Mathematics. 6(6) (2021), 5518-5534.
[8] G A. Derfel, A. Iserles, The pantograph equation in the complex plane, J Math Anal Appl. 213, (1997), 117-132.
[9] M. Houas, Existence of solutions for a coupled system of Caputo-Hadamard type fractional differential equations with Hadamard fractional integral conditions, Adv.Theory Nonlinear Anal. Appl. 5(3) (2021), 316-329.
[10] A. Iserles, Exact and discretized stability of the pantograph equation, Appl. Numer. Math. 24 (1997), 295-308.
[11] A Iserles, Y. Liu, On pantograph integro-differential equations, J. Integral Equations Appl. 6 (1994), 213-237.
[12] A. Khan, H. Khan, J.F.Gómez-Aguilar and T. Abdeljawad, Existence and Hyers-Ulam stability for a nonlinear singular fractional differential equations with Mittag-Leffler kernel, Chaos Solitons Fractals. 127 (2019), 422-427.
[13] A.Khan, J.F. Gómez-Aguilar, T. Abdeljawada and H. Khand, Stability and numerical simulation of a fractional order plant-nectar-pollinator model, Alexandria Engineering Journal. 59 (2020), 49-59.
[14] A. Khan, Y. Li, K. Shah and T. S. Khan, On coupled $p-$Laplacian fractional differential equations with nonlinear boundary conditions, Complexity. 2017, Article ID 8197610: (2017), 1-9.
[15] H. Khan, W. Chen, A. Khan, T. S. Khan and Q. M. Al-Madlal, Hyers-Ulam stability and existence criteria for coupled fractional differential equations involving $p-$Laplacian operator, Adv. Difference . Equ. 2018: 45 (2018), 1-16.
[16] H. Khan, C. Tunc and A. Khan, Green function's properties and existence theorems for nonlinear singular-delay-fractional differential equations, Discrete Contin. Dyn. Syst., Ser. S. 13(9) (2020), 2475-2487.
[17] H. Khan, J.F.Gomez-Aguilar, T. Abdeljawad and A. Khan, Existence results and stability criteria for abc-fuzzy-Volterra integro-differential equation, Fractals. 28(8) (2020), 204004-1-204004-9.
[18] M. B. A. Khan, T. Abdeljawad, K. Shah, G. Ali, H. Khan and A. Khan, Study of a nonlinear multi-terms boundary value problem of fractional pantograph differential equations, Adv. Difference Equ. 2021: 143 (2021), 1-15.
[19] A.A. Kilbas, S.A. Marzan. Nonlinear differential equation with the Caputo fraction derivative in the space of continuously differentiable functions. Differ. Equ. 41(1), (2005), 84-89.
[20] M. Klimek, Sequential fractional differential equations with Hadamard derivative, Commun. Nonlinear Sci. Numer. Simul. 16 (2011), 4689-4697.
[21] V. Lakshmikantham, S.Leela, and J. V.Devi. Theory of fractional dynamic systems. Cambridge Scientific Publishers. 2009.
[22] N. Mahmudov, M. Awadalla and K. Abuassba, Nonlinear sequential fractional differential equations with nonlocal boundary conditions, Adv. Difference Equ. 2017: 319 (2017), 1-.15
[23] J R. Ockendon, A B. Taylor, *The dynamics of a current collection system for an electric locomotive*, Proc RSoc London, Ser.A. 322 (1971), 447-468.
[24] A. Seemab, M. U. Rehman, J. Alzabut, Y. Adjabi and M.S. Abdo, Langevin equation with nonlocal boundary conditions involving a $\psi-$Caputo fractional operators of different orders, AIMS Mathematics. 6(7) (2021), 6749-6780.
[25] D. Vivek, E. M. Elsayed and K. Kanagarajan, Existence and Ulam stability results for a class of boundary value problem of neutral pantograph equations with complex order, SeMA Journal. 77(3) (2021), 243-256..
[26] H. A. Wahashy, M.S. Abdo, A. M. Saeed and S.K. Panchal, Singular fractional differential equations with $\psi-$Caputo operator and modified Picard's iterative method. Appl. Math. E-Notes. 20 (2020), 215-229.
[27] Z. Wei, W. Dong, Periodic boundary value problems for Riemann-Liouville sequential fractional differential equations, Electron. J. Qual. Theory Differ. Equ. 87 (2011), 1-13.
[28] A. Wongcharoen, S.K. Ntouyas and J.Tariboon, Nonlocal boundary value problemsfor Hilfer-type pantograph fractionald-ifferential equations and inclusions, Adv. Difference. Equ. 2020: 279 (2020), 1-21.
[29] A. Zada, M. Yar and T. Li, Existence and stability analysis of nonlinear sequential coupled system of Caputo fractional differential equations with integral boundary conditions, Ann. Univ. Paedagog. Crac. Stud. Math. 17 (2018), 103-125.