



Several recurrence relations and identities on generalized derangement numbers

Muhammet Cihat Dağlı^a, Feng Qi^{b,c,d}

^aDepartment of Mathematics, Akdeniz University, 07058-Antalya, Turkey

^bInstitute of Mathematics, Henan Polytechnic University, Jiaozuo 454010, Henan, China

^cSchool of Mathematical Sciences, Tiangong University, Tianjin 300387, China

^dCorresponding author

Abstract

In the paper, with aid of generating functions, the authors present several recurrence relations and identities for generalized derangement numbers involving generalized harmonic numbers and the Stirling numbers of the first kind.

Keywords: recurrence relation; identity; generalized derangement number; generalized harmonic numbers; Stirling numbers of the first kind; generating function

2010 MSC: 05A15; 05A19; 11B73; 11B83; 11B37; 11Y55

1. Introduction

In combinatorial mathematics, a derangement is a permutation of the elements of a set, such that no element appears in its original position. The number of derangements of a set of size n is called the derangement number. Derangement numbers can be generated by

$$\frac{e^{-t}}{1-t} = \sum_{n=0}^{\infty} d_n \frac{t^n}{n!}$$

and possess the closed-form expression

$$d_n = \sum_{k=0}^n (-1)^k \frac{n!}{k!}.$$

Email addresses: mcihatdagli@akdeniz.edu.tr (Muhammet Cihat Dağlı), qifeng618@gmail.com, qifeng618@hotmail.com, qifeng618@qq.com (Feng Qi)

Received September 29, 2021; Accepted: April 24, 2022; Online: April 28, 2022

See the monograph [3] and the papers [18, 20, 21, 23, 24, 25, 26].

The generalized derangement numbers $d_{n,r}$ are introduced by Munarini [15] as

$$d_{n,r} = \sum_{k=0}^n (-1)^k \binom{r+n-k}{n-k} \frac{n!}{k!}$$

and can be generated by

$$\frac{e^{-t}}{(1-t)^{r+1}} = \sum_{n=0}^{\infty} d_{n,r} \frac{t^n}{n!}. \quad (1)$$

It is clear that $d_{n,0} = d_n$. The first few generalized derangement numbers $d_{n,r}$ are

$$d_{0,r} = 1, \quad d_{1,r} = r, \quad d_{2,r} = r^2 + r + 1, \quad d_{3,r} = r^3 + 3r^2 + 5r + 2.$$

Several types of extensions of derangement numbers have been offered and many of remarkable properties and identities have been achieved for them. See, for example, the papers [12, 13, 14, 30, 31, 32].

Harmonic numbers [9] are located in a very important position in combinatorial number theory. Various generalizations for these numbers have been introduced in the literature. One of them, due to [5], is denoted by $H(n, r, \alpha)$ and defined by

$$\sum_{n=0}^{\infty} H(n, r, \alpha) t^n = \frac{[-\ln(1 - \frac{t}{\alpha})]^{r+1}}{1-t}. \quad (2)$$

Employing generating functions, the authors of [5] established a number of combinatorial identities and relations for generalized harmonic numbers $H(n, r, \alpha)$ and some important special polynomials and numbers. Setting $r = 0$ in (2) leads to another type of generalized harmonic numbers, which are given by

$$H_0(\alpha) = 0 \quad \text{and} \quad H_n(\alpha) = \sum_{i=1}^n \frac{1}{i\alpha^i}$$

for $n \in \mathbb{N} = \{1, 2, \dots\}$ and can be generated by

$$\sum_{n=1}^{\infty} H_n(\alpha) t^n = \frac{-\ln(1 - \frac{t}{\alpha})}{1-t}.$$

Note that $H_n(1) = H_n$ denotes the classical harmonic numbers [6, 16]. Furthermore, a generalization for hyperharmonic numbers was presented in [17] as

$$H_{n,r}(\alpha) = \sum_{i=1}^n H_{i,r-1}(\alpha), \quad r, n \geq 1$$

subject to initial condition $H_{n,0}(\alpha) = \frac{1}{n\alpha^n}$, and can be generated by

$$\sum_{n=0}^{\infty} H_{n,r}(\alpha) t^n = \frac{-\ln(1 - \frac{t}{\alpha})}{(1-t)^r}. \quad (3)$$

For further investigations concerning with generalized harmonic numbers, the readers may consult with [1, 2, 4, 7, 10, 11, 27, 28] and references cited therein.

As another important number in combinatorial analysis, the Stirling numbers of the first kind can be generated by

$$\frac{[\ln(1+x)]^k}{k!} = \sum_{n=k}^{\infty} s(n, k) \frac{x^n}{n!}, \quad |x| < 1 \quad (4)$$

and satisfy the (diagonal) recurrence relations

$$\begin{aligned}
 s(n + 1, k) &= s(n, k - 1) - ns(n, k), \\
 s(n, k) &= (-1)^k \sum_{m=1}^n (-1)^m \sum_{\ell=k-m}^{k-1} (-1)^\ell \binom{n}{\ell} \binom{\ell}{k-m} s(n - \ell, k - \ell) \\
 &= (-1)^{n-k} \sum_{\ell=0}^{k-1} (-1)^\ell \binom{n}{\ell} \binom{\ell-1}{k-n-1} s(n - \ell, k - \ell),
 \end{aligned}$$

and

$$\frac{s(n + k, k)}{\binom{n+k}{k}} = \sum_{\ell=0}^n (-1)^\ell \frac{\langle k \rangle_\ell}{\ell!} \sum_{m=0}^{\ell} (-1)^m \binom{\ell}{m} \frac{s(n + m, m)}{\binom{n+m}{m}}, \tag{5}$$

where the conventions that $\binom{0}{0} = 1$, $\binom{-1}{-1} = 1$, and $\binom{p}{q} = 0$ for $p \geq 0 > q$ are adopted and

$$\langle x \rangle_n = \prod_{k=0}^{n-1} (x - k) = \begin{cases} x(x - 1) \cdots (x - n + 1), & n \geq 1 \\ 1, & n = 0 \end{cases}$$

is the falling factorial. See the book [8] and the papers [19, 22].

The classical Euler’s gamma function $\Gamma(z)$ can be defined [29, Chapter 3] by

$$\Gamma(z) = \lim_{n \rightarrow +\infty} \frac{n! n^z}{\prod_{k=0}^n (z + k)}, \quad z \in \mathbb{C} \setminus \{0, -1, -2, \dots\}.$$

The extended binomial coefficient $\binom{z}{w}$ for $z, w \in \mathbb{C}$ is defined by

$$\binom{z}{w} = \begin{cases} \frac{\Gamma(z + 1)}{\Gamma(w + 1)\Gamma(z - w + 1)}, & z \notin \mathbb{N}_-, \quad w, z - w \notin \mathbb{N}_-; \\ 0, & z \notin \mathbb{N}_-, \quad w \in \mathbb{N}_- \text{ or } z - w \in \mathbb{N}_-; \\ \frac{\langle z \rangle_w}{w!}, & z \in \mathbb{N}_-, \quad w \in \mathbb{N}_0; \\ \frac{\langle z \rangle_{z-w}}{(z - w)!}, & z, w \in \mathbb{N}_-, \quad z - w \in \mathbb{N}_0; \\ 0, & z, w \in \mathbb{N}_-, \quad z - w \in \mathbb{N}_-; \\ \pm\infty, & z \in \mathbb{N}_-, \quad w \notin \mathbb{Z}; \end{cases}$$

where

$$\begin{aligned}
 \mathbb{R} &= (-\infty, \infty), & \mathbb{C} &= \{x + iy : x, y \in \mathbb{R}, i = \sqrt{-1}\}, & \mathbb{Z} &= \{0, \pm 1, \pm 2, \dots\}, \\
 \mathbb{N} &= \{1, 2, \dots\}, & \mathbb{N}_0 &= \{0, 1, 2, \dots\}, & \mathbb{N}_- &= \{-1, -2, \dots\}.
 \end{aligned}$$

In this paper, with help of generating functions, we will present several recurrence relations and identities for generalized derangement numbers $d_{n,r}$ involving generalized harmonic numbers $H(k, r, \alpha)$ and the Stirling numbers of the first kind $s(n, k)$.

2. Main results and their proofs

This section is devoted to demonstrate our main results and their proofs.

Theorem 1. *The generalized derangement numbers $d_{n,r}$ satisfy*

$$\sum_{k=0}^n \binom{n}{k} d_{k,r} = n! \binom{n+r}{n}, \quad n \in \mathbb{N}_0.$$

Proof. From the generating function in (1), it is ready that

$$\frac{1}{(1-t)^{r+1}} = e^t \sum_{n=0}^{\infty} d_{n,r} \frac{t^n}{n!}.$$

Using the binomial theorem and applying the Cauchy product, we have

$$\sum_{n=0}^{\infty} \binom{-r-1}{n} (-t)^n = \left(\sum_{n=0}^{\infty} \frac{t^n}{n!} \right) \sum_{n=0}^{\infty} d_{n,r} \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left[\sum_{k=0}^n \frac{d_{n-k,r}}{k!(n-k)!} \right] t^n. \tag{6}$$

Comparing the coefficients of t^n on both sides of (6) and using the identity

$$\binom{x}{k} = (-1)^k \binom{-x+k-1}{k}$$

complete the proof. □

Remark 1. For $r = 0$, Theorem 1 coincides with the second equality in [26, Theorem 2].

Theorem 2. Generalized derangement numbers $d_{n,r}$ satisfy

$$d_{n,r+1} = d_{n,r} + nd_{n-1,r+1}. \tag{7}$$

Proof. From the equation (1), we have

$$\sum_{n=0}^{\infty} (d_{n,r+1} - d_{n,r}) \frac{t^n}{n!} = \frac{e^{-t}}{(1-t)^{r+2}} - \frac{e^{-t}}{(1-t)^{r+1}} = t \frac{e^{-t}}{(1-t)^{r+2}} = \sum_{n=0}^{\infty} d_{n,r+1} \frac{t^{n+1}}{n!} = \sum_{n=0}^{\infty} nd_{n-1,r+1} \frac{t^n}{n!}.$$

Equating the coefficients $\frac{t^n}{n!}$ gives the desired result. □

Theorem 3. The generalized derangement numbers $d_{n,r}$ satisfy the recurrence relations

$$d_{n+1,r} = rd_{n,r+1} + nd_{n-1,r+1} \tag{8}$$

and

$$d_{n+1,r} = (r+1)d_{n,r+1} - d_{n,r}. \tag{9}$$

Proof. From the generating function in (1), we acquire

$$\frac{d}{dt} \left[\frac{e^{-t}}{(1-t)^{r+1}} \right] = \sum_{n=0}^{\infty} d_{n+1,r} \frac{t^n}{n!}. \tag{10}$$

Differentiating on both sides of the equation (1) with respect to t results in

$$\frac{d}{dt} \left[\frac{e^{-t}}{(1-t)^{r+1}} \right] = (r+t) \frac{e^{-t}}{(1-t)^{r+2}} = (r+t) \sum_{n=0}^{\infty} d_{n,r+1} \frac{t^n}{n!} = r \sum_{n=0}^{\infty} d_{n,r+1} \frac{t^n}{n!} + \sum_{n=0}^{\infty} nd_{n-1,r+1} \frac{t^n}{n!}. \tag{11}$$

Combining (10) and (11) lead to the equation (8).

If subtracting (7) from (8), we obtain the relation (9) immediately. □

Theorem 4. Let $n, r \in \mathbb{N}$. Then

$$\sum_{k=0}^n \frac{(-1)^{n-k}}{(n-k)!} H_{k,r+1}(\alpha) = \sum_{k=0}^n \frac{d_{n-k,r-1}}{(n-k)!} H_k(\alpha).$$

Proof. From the generating function in (3), one can write

$$\sum_{n=0}^{\infty} H_{n,r+1}(\alpha)t^n = \frac{-\ln(1-\frac{t}{\alpha})}{(1-t)^{r+1}} = \frac{-\ln(1-\frac{t}{\alpha})}{1-t} \frac{e^{-t}}{(1-t)^r} \frac{1}{e^{-t}},$$

from which it is easy to verify that

$$\sum_{n=0}^{\infty} H_{n,r+1}(\alpha)t^n \sum_{n=0}^{\infty} \frac{(-t)^n}{n!} = \sum_{n=1}^{\infty} H_n(\alpha)t^n \sum_{n=0}^{\infty} d_{n,r-1} \frac{t^n}{n!}.$$

Applying the Cauchy product and equating the coefficients of t^n yield the desired formula. \square

Theorem 5. Let $n, r \in \mathbb{N}$. Then

$$\sum_{k=0}^n k!H(k, r, \alpha)d_{n-k,r-1} = (-1)^{n-r-1} \frac{(r+1)!}{\alpha^n} \sum_{k=0}^n (-\alpha)^k s(n-k, r+1)d_{k,r}.$$

Proof. Consider the equality

$$\frac{[-\ln(1-\frac{t}{\alpha})]^{r+1}}{1-t} \frac{e^{-t}}{(1-t)^r} = \left[-\ln\left(1-\frac{t}{\alpha}\right)\right]^{r+1} \frac{e^{-t}}{(1-t)^{r+1}}.$$

Utilizing the generating functions in (1), (2), and (4), we find

$$\sum_{n=0}^{\infty} H(n, r, \alpha)t^n \sum_{n=0}^{\infty} d_{n,r-1} \frac{t^n}{n!} = \sum_{n=0}^{\infty} \frac{(-1)^{n-r-1}(r+1)!s(n, r+1)}{\alpha^n} \frac{t^n}{n!} \sum_{n=0}^{\infty} d_{n,r} \frac{t^n}{n!},$$

from which, we have

$$\sum_{n=0}^{\infty} \left[\sum_{k=0}^n k!H(k, r, \alpha)d_{n-k,r-1} \right] \frac{t^n}{n!} = \sum_{n=0}^{\infty} \left[\sum_{k=0}^n d_{k,r} \frac{(-1)^{n-k-r-1}(r+1)!s(n-k, r+1)}{\alpha^{n-k}} \right] \frac{t^n}{n!}.$$

Accordingly the desired relation follows immediately. \square

References

- [1] G.-S. Cheon and M. El-Mikkawy, *Generalized harmonic numbers with Riordan arrays*, J. Number Theory **128** (2008), no. 2, 413–425; available online at <https://doi.org/10.1016/j.jnt.2007.08.011>.
- [2] J. Choi and H. M. Srivastava, *Some summation formulas involving harmonic numbers and generalized harmonic numbers*, Math. Comput. Modelling **54** (2011), 2220–2234; available online at <https://doi.org/10.1016/j.mcm.2011.05.032>.
- [3] L. Comtet, *Advanced Combinatorics: The Art of Finite and Infinite Expansions*, Revised and Enlarged Edition, D. Reidel Publishing Co., 1974; available online at <https://doi.org/10.1007/978-94-010-2196-8>.
- [4] G. Dattoli, S. Licciardi, E. Sabia, and H. M. Srivastava, *Some properties and generating functions of generalized harmonic numbers*, Mathematics **7** (2019), no. 4, Article No. 577; available online at <https://doi.org/10.3390/math7070577>.
- [5] Ö. Duran, N. Ömür, and S. Koparal, *On sums with generalized harmonic, hyperharmonic and special numbers*, Miskolc Math. Notes **21** (2020), no. 2, 791–803; available online at <https://doi.org/10.18514/MMN.2020.3458>.
- [6] C.-J. Feng and F.-Z. Zhao, *Some results for generalized harmonic numbers*, Integers **9** (2009), no. 5, 605–619; available online at <https://doi.org/10.1515/INTEG.2009.048>.
- [7] A. Gertsch, *Nombres harmoniques généralisés*, C. R. Acad. Sci. Paris Sér. I Math. **324** (1997), no. 1, 7–10; available online at [https://doi.org/10.1016/S0764-4442\(97\)80094-8](https://doi.org/10.1016/S0764-4442(97)80094-8). (French)
- [8] R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete Mathematics: A Foundation for Computer Science*, Addison-Wesley Publishing Company, Reading, MA, 1994.
- [9] B.-N. Guo and F. Qi, *Sharp inequalities for the psi function and harmonic numbers*, Analysis (Berlin) **34** (2014), no. 2, 201–208; available online at <https://doi.org/10.1515/anly-2014-0001>.
- [10] L. Kargin and M. Can, *Harmonic number identities via polynomials with r -Lah coefficients*, C. R. Math. Acad. Sci. Paris **358** (2020), no. 5, 535–550; available online at <https://doi.org/10.5802/crmath.53>.

- [11] L. Kargin, M. Cenkci, A. Dil, and M. Can, *Generalized harmonic numbers via poly-Bernoulli polynomials*, arXiv (2008), available online at <https://arxiv.org/abs/2008.00284v2>.
- [12] T. Kim and D. S. Kim, *Some identities on derangement and degenerate derangement polynomials*, Advances in Mathematical Inequalities and Applications, 265–277, Trends Math., Birkhäuser/Springer, Singapore, 2018; available online at https://doi.org/10.1007/978-981-13-3013-1_13.
- [13] T. Kim, D. S. Kim, D. V. Dolgy, and J. Kwon, *Some identities of derangement numbers*, Proc. Jangjeon Math. Soc. **21** (2018), no. 1, 125–141.
- [14] T. Kim, D. S. Kim, G.-W. Jang, and J. Kwon, *A note on some identities of derangement polynomials*, J. Inequal. Appl. **2018**, Paper No. 40, 17 pages; available online at <https://doi.org/10.1186/s13660-018-1636-8>.
- [15] E. Munarini, *Callan-like identities*, Online J. Anal. Comb. No. 14 (2019), 20 pp.
- [16] D.-W. Niu, Y.-J. Zhang, and F. Qi, *A double inequality for the harmonic number in terms of the hyperbolic cosine*, Turkish J. Anal. Number Theory **2** (2014), no. 6, 223–225; available online at <http://dx.doi.org/10.12691/tjant-2-6-6>.
- [17] N. Ömür and G. Bilgin, *Some applications of the generalized hyperharmonic numbers of order r , $H_n^r(\alpha)$* , Adv. Appl. Math. Sci. **17** (2018), no. 9, 617–627.
- [18] F. Qi, *A determinantal representation for derangement numbers*, Glob. J. Math. Anal. **4** (2016), no. 3, 17–17; available online at <https://doi.org/10.14419/gjma.v4i3.6574>.
- [19] F. Qi, *Diagonal recurrence relations for the Stirling numbers of the first kind*, Contrib. Discrete Math. **11** (2016), 22–30; available online at <https://doi.org/10.11575/cdm.v11i1.62389>.
- [20] F. Qi, *Three closed forms for convolved Fibonacci numbers*, Results Nonlinear Anal. **3** (2020), no. 4, 185–195.
- [21] F. Qi, M. C. Dağlı, and W.-S. Du, *Determinantal forms and recursive relations of the Delannoy two-functional sequence*, Adv. Theory Nonlinear Anal. Appl. **4** (2020), no. 3, 184–193; available online at <https://doi.org/10.31197/atnaa.772734>.
- [22] F. Qi and B.-N. Guo, *A diagonal recurrence relation for the Stirling numbers of the first kind*, Appl. Anal. Discrete Math. **12** (2018), no. 1, 153–165; available online at <https://doi.org/10.2298/AADM170405004Q>.
- [23] F. Qi and B.-N. Guo, *Explicit formulas for derangement numbers and their generating function*, J. Nonlinear Funct. Anal. **2016**, Article ID 45, 10 pages.
- [24] F. Qi, J.-L. Wang, and B.-N. Guo, *A recovery of two determinantal representations for derangement numbers*, Cogent Math. (2016), **3**:1232878, 7 pages; available online at <https://doi.org/10.1080/23311835.2016.1232878>.
- [25] F. Qi, J.-L. Wang, and B.-N. Guo, *A representation for derangement numbers in terms of a tridiagonal determinant*, Kragujevac J. Math. **42** (2018), no. 1, 7–14; available online at <https://doi.org/10.5937/KgJMath1801007F>.
- [26] F. Qi, J.-L. Zhao, and B.-N. Guo, *Closed forms for derangement numbers in terms of the Hessenberg determinants*, Rev. R. Acad. Cienc. Exactas Fis. Nat. Ser. A Mat. RACSAM **112** (2018), no. 4, 933–944; available online at <https://doi.org/10.1007/s13398-017-0401-z>.
- [27] J. M. Santmyer, *A Stirling like sequence of rational numbers*, Discret. Math. **171** (1997), no. 3, 229–235; available online at [https://doi.org/10.1016/S0012-365X\(96\)00082-9](https://doi.org/10.1016/S0012-365X(96)00082-9).
- [28] A. Sofo and H. M. Srivastava, *A family of shifted harmonic sums*, Ramanujan J. **37** (2015), no. 1, 89–108; available online at <https://doi.org/10.1007/s11139-014-9600-9>.
- [29] N. M. Temme, *Special Functions: An Introduction to Classical Functions of Mathematical Physics*, A Wiley-Interscience Publication, John Wiley & Sons, Inc., New York, 1996; available online at <http://dx.doi.org/10.1002/9781118032572>.
- [30] M. L. Wachs, *On q -derangement numbers*, Proc. Amer. Math. Soc. **106** (1989), no. 1, 273–278; available online at <https://doi.org/10.2307/2047402>.
- [31] C. Y. Wang, P. Miska, and I. Mezö, *The r -derangement numbers*, Discret. Math. **340** (2017), 1681–1692; available online at <https://doi.org/10.1016/j.disc.2016.10.012>.
- [32] F.-Z. Zhao, *The log-balancedness of generalized derangement numbers*, Ramanujan J. (2021), in press; available online at <https://doi.org/10.1007/s11139-021-00402-1>.